

**ANIMVET COMPANY
PRODUCTS
CATALOGUE**

HAYAT ROKHAN LTD

PRODUCTS:	PAGE:
ANIPARA C WS.....	4
ANIELECTROLYTE WS.....	5
ANISTRES WS.....	6
ANIVIT EC WS.....	7
ANINTITOX WS.....	8
ANIGUMBURO WS.....	9
ANIVIT EC WS.....	10
ANIMINOMIX PLUS WS.....	11
ANIVIT A+.....	13
ANIGARLIMINT SUPER.....	14
ANITONIC.....	15
ANIMIN.....	16
ANIMIN SUPER.....	17
ANIRES P.....	18
ANIVIT AD3ECK.....	19
ANIVIT AMINO.....	20
ANIVIT CARLYMES.....	21

HAYAT ROKHAN LTD



ANIMVET COMPANY

PRODUCTS

Water Soluble

Powders

HAYAT ROKHAN LTD

ANIPARA C WS

Powder for oral administration

PRODUCT CATEGORY

VETERINARY PHARMACEUTICAL

THERAPEUTIC GROUP

ANALGESIC

PHARMACOTHERAPEUTIC GROUP

Analgesics and antipyretics

COMPOSITION

Contains per gram powder:

Paracetamol.....	250 mg/g
Ascorbic acid.....	250 mg/g
Methionine.....	100 mg/g

Withdrawal Times:

None

Storage:

Dark, dry and bellow 25 C. Do not freeze.

TARGET SPECIES

Poultry, pig

PRESENTATION

1 kg

INDICATIONS

Symptomatic treatment of fever in the context of respiratory diseases in combination with an appropriate anti-infective therapy, if necessary.

DOSAGE

Oral route

Poultry: 250-500 g/1000 L drinking water, for 3-5days

Pigs: 15-30 mg of paracetamol/kg bw/day, as long as the pigs are suffering from pyrexia for a maximum duration treatment of 5 days (equivalent to 1.5 g of oral powder per 10 kg bodyweight).

WITHDRAWAL PERIOD

Meat and offal: zero days.

ANIELECTROLYTE WS

Powder for oral administration

PRODUCT CATEGORY:

Feed Additive

THERAPEUTIC GROUP:

Nutritional

PHARMACOTHERAPEUTIC GROUP:

Vitamin Combination

COMPOSITION:

Sodium chloride.....110 g/kg
Potassium chloride.....13.5 g/kg
Sodium dihydrogen phosphate.....22 g/kg
Sodium citrate.....5 g/kg
Glycine.....45 g/kg
Dextrose monohydrate q.s.p.

Withdrawal Times:

None

Storage:

Dark, dry and bellow 25 C. Do not freeze.

TARGET SPECIES:

Poultry

DIRECTIONS FOR USE:

ANIELECTROL YTE powder is using as a supportive in the treatment of metabolic diseases due to electrolyte deficiency (dehydration, temperature stress etc.) and acid base imbalance cases. It can also be used as a carbohydrate source.

DOSAGE:

Poultry: 500 g-1 kg/ton feed or drinking water, for 3-5 days.

PRESENTATION: 1 Kg

HAYAT ROKHAN LTD

ANISTRES WS

Powder for oral administration

PRODUCT CATEGORY:

Feed Additive

THERAPEUTIC GROUP:

Nutritional

PHARMACOTHERAPEUTIC GROUP:

Vitamin-Mineral

COMPOSITION:

Sodium chloride.....	200 g/kg
Potassium chloride.....	200 g/kg
Calcium sulphate.....	100 g/kg
Vitamin C.....	50 g/kg
Vitamin B2.....	910 mg/kg
Vitamin B3.....	3500 mg/kg
Vitamin B6.....	1220 mg/kg
Vitamin B12.....	2260 mg/kg

Withdrawal Times:

None

Storage:

Dark, dry and bellow 25 C. Do not freeze.

TARGET SPECIES:

Poultry, ruminant, horse

DIRECTIONS FOR USE:

ANISTRES WS powder used as a supplement to meet the daily needs of electrolytes, vitamin B complex and vitamin C in poultry, cattle and horses. Recommended to be used for supporting the conditions of stress caused by vaccinations, diseases, transport, high humidity, high temperatures or extreme temperature changes. It can also be used for improving body growth, feed consumption rate, immune system, productivity and egg cortex quality. It supplies the strength and elasticity of vein and arteries, helps to prevent acidosis and alkalosis.

DOSAGE:

Poultry: 1 kg/1 ton drinking water or feed

Cattle and camels (not-training): 1 kg/1 ton drinking water or feed

Horses (not-training): 50-60 g/animal

PRESENTATION: 1 kg

ANIVIT EC WS

Powder for oral administration

PRODUCT CATEGORY:

Feed Additive

THERAPEUTIC GROUP:

Nutritional

PHARMACOTHERAPEUTIC GROUP:

Vitamin-Mineral

COMPOSITION:

Sodium selenite.....450 mg/L

Vitamin E.....100000 mg/L

Withdrawal Times:

None

Storage:

Dark, dry and bellow 25 C. Do not freeze.

TARGET SPECIES:

Poultry, pig

DIRECTIONS FOR USE:

ANIVIT EC Powder for oral solution is used to support for

- improvement of fertility, feed conversion rate and carcass quality, egg and hatch efficiency,
- protective treatment of muscle dystrophy, pancreas encephalomalacia (epidemic tremor), exudative diathesis and myopathy.
- prevention and treatment of toxication,
- stimulation of the immune system after vaccination.

DOSAGE:

Poultry: 1 L/ 1000 L drinking water/day Sheep, lambs and pigs: 5-10 mL/animal

PRESENTATION: 1 KG

HAYAT ROKHAN LTD

ANINTITOX WS

Powder for oral administration

PRODUCT CATEGORY:

Feed Additive

THERAPEUTIC GROUP:

Nutritional

PHARMACOTHERAPEUTIC GROUP:

Vitamin-Mineral

COMPOSITION:

Vitamin A E.....2.5 g/kg
Vitamin C.....2.5 g/kg
Vitamin K.....1 g/kg
Sodiumbenzoate.....500 g/kg
Sodium dihydrogen phosphate.....100 g/kg

Withdrawal Times:

None

Storage:

Dark, dry and bellow 25 C. Do not freeze.

TARGET SPECIES:

Poultry

Indications:

ANINTITOX is a multivitamin and mineral feed additive composition used for supplying the vitamin A, vitamin E, vitamin C, vitamin K3 and sodium deficiencies to correct concurrent clinical or subclinical deficiencies of poultry. Strategic vitamin supplementation is aimed at the optimization of the gastro-intestinal and reproductive performance, breeding life cycle, immune function and animal growth. It can also be used as tonic for liver and kidneys.

DOSAGE:

Poultry: 500 g-1 kg /1ton feed or drinking water

PRESENTATION: 1 Kg

HAYAT ROKHAN LTD

ANIGUMBURO WS

Powder for oral administration

PRODUCT CATEGORY:

Feed Additive

THERAPEUTIC GROUP:

Nutritional

PHARMACOTHERAPEUTIC GROUP:

Vitamin & Amino acid & Systemic-urinary acidifying salt

COMPOSITION:

Ammonium chloride.....	700 g/kg
Methionine.....	40 g/kg
Vitamin A.....	1000000 IU/kg
Vitamin E.....	1 g/kg
Vitamin C.....	40 g/kg
Vitamin B1.....	500 mg/kg
Vitamin B2.....	400 mg/kg
Vitamin B6.....	400 mg/kg
Vitamin K3.....	2 g/kg
Dextrose.....	50 g/kg

Withdrawal Times:

None

Storage:

Dark, dry and bellow 25 C. Do not freeze.

TARGET SPECIES:

Poultry, ruminant

DIRECTIONS FOR USE:

ANIGUMBURO is a feed additive combination of vitamin A, vitamin E, vitamin C, B group of vitamins, vitamin K3, ammonium chloride and methionine which can be used In vitamin deficiency cases. Recommended to be used as a supportive as diuretic, liver & kidney detoxificant and immune booster.

DOSAGE:

Poultry: 500 g/ton drinking water for 3-5 days

PRESENTATION: 1 Kg

HAYAT ROKHAN LTD

ANIVIT EC WS

Powder for oral administration

PRODUCT CATEGORY:

Feed Additive

THERAPEUTIC GROUP:

Nutritional

PHARMACOTHERAPEUTIC GROUP:

Vitamin-Mineral

COMPOSITION:

Vitamin A.....	15.000.000 IU/kg
Vitamin D3.....	1.500.000 IU/kg
Vitamin E.....	2000 IU/kg
Vitamin B1.....	2000 mg/kg
Vitamin B2.....	2500 mg/kg
Vitamin B6.....	3000 mg/kg
Vitamin B12.....	5 mg/kg
Nicotinamide.....	10.000 mg/kg
Calcium pantothenate.....	5500 mg/kg
Vitamin K.....	2000 mg/kg
Folic acid.....	2 mg/kg

Withdrawal Times:

None

Storage:

Dark, dry and bellow 25 C. Do not freeze.

TARGET SPECIES:

Poultry, ruminant, horse, pig

DIRECTIONS FOR USE:

ANIVIT ECO Powder for oral solution is used in deficiency cases of vitamin A, D3, E, B group vitamins (B1, B2, B3, B6, B9, B12), vitamin K and calcium in its composition. It is used to support for improvement of

- immune system
- body resistance in the conditions of immunosuppression and in the recovery periods of diseases and stress conditions
- quality of meat and egg

DOSAGE:

Poultry: 200 g-300 g/ 1000 L drinking water/day

Cattle, camels and horses: 10 g animal/day Calves, foals and pigs: 5 g animal/day

PRESENTATION: 1 Kg

ANIMINOMIX PLUS WS

Powder for oral administration

PRODUCT CATEGORY:

Feed Additive

THERAPEUTIC GROUP:

Nutritional

PHARMACOTHERAPEUTIC GROUP:

Vitamin & Minerals

Composition:

26000 IU/kgVitamin A
52000 IU/kgVitamin D3
120 mg/kgVitamin E
500 mg/kgVitamin B3
2500 mg/kgIron (Fe)
80 mg/kgCalcium iodate
10 mg/kgCobalt chloride
500 mg/kgCopper sulfate
2500 mg/kgManganese chloride
2500 mg/kgZinc sulfate
10 mg/kgSodium selenite 3000
mg/kgDicalcium phosphate
18000 mg/kgSodium chloride
9000 mg/kgMagnesium sulfate
200 mg/kgL-Arginine
200 mg/kgL-Lysine
200 mg/kgDL-Methionine
6.6 x10⁸ CFU/kgYeast (*Saccharomyces cerevisiae*) 1530 mg/kg Inactive yeast
product (*Saccharomyces cerevisiae*) Orange flavor Calcium carbonate q.s.p.

Withdrawal Times:

None

Storage:

Dark, dry and bellow 25 C. Do not freeze.

Benefits:

Supportive during stress Boosts the immune system Contribute to an optimal function of the reproductive system Improve egg and meat quality Stimulate muscle growth Contributes to optimal milk production Recommended for animal in high demanding –production stages Simple administration through drinking water

Dosage:

For oral administration through drinking water.

Cattle: 1g per 40 kg body weight for 3-5 days

Calves, sheep and goats: 1g per 20 kg body weight for 3-5 days

Swine and Poultry: 1 kg per 2000 liters of drinking water for 3-5 days

PRESENTATION: 1 Kg

**ANIMVIT COMPANY
PRODUCTS
Oral Solutions**

HAYAT ROKHAN LTD

ANIVIT A+

Solution for Oral

PRODUCT CATEGORY:

Feed Additive

THERAPEUTIC GROUP:

Nutritional

PHARMACOTHERAPEUTIC GROUP:

Vitamin & Amino acid

Composition:

8 000 000 IU/L	Vitamin A	300 mg /L	Biotin
800 000 IU/L	Vitamin D3	1200 mg/L	L-lysine monohydrochloride
4000 mg/L	Vitamin E	200 mg/L	DL-methionine
10000 mg/L	Vitamin C	200 mg/L	L-threonine
1000 mg/L	Vitamin B1	8 mg/L	L-tryptophan
200 mg/L	Vitamin B2	200 mg/L	L-valine
200 mg/L	Vitamin B6	200 mg/L	L-arginine
200 mg/L	Vitamin B12	200 mg/L	L-leucin
400 mg/L	Vitamin K3	80 mg/L	L-isoleucine
28000 mg/L	Choline chloride	80 mg/L.....	L-tyrosine
500 mg/L	Calcium-DPanthotenat	300 mg/L	L-Proline
1600 mg/L	L-Glutamic Acid		

Withdrawal Times:

None

Storage:

Dark, dry and bellow 25 C. Do not freeze.

TARGET SPECIES:

Poultry, ruminant, horse, pet, pig

DIRECTIONS FOR USE:

ANIVIT A+ oral Solution for oral recommended using as a supportive supplement in cases of dehydration, electrolyte, and vitamin imbalance, hyperproteinemia that can occur in diarrhea, shock and exhaustion. Recommended using as a preventative supplement in cases of breeding losses, nutritional deficiencies cases, and fatty liver syndrome and nerve disorders. In addition, it is using during period of convalescence.

DOSAGE:

Poultry: 250-500 mL/ton drinking water, for 3-5 days
Horse: 20-30 mL/day Calves: 10-20 mL/day
Pigs: 5-10 mL Lambs: 2-4 mL/day Cats and dogs: 1-2 mL/day

PRESENTATION: 1 L

ANIGARLIMINT SUPER

Solution for Oral

PRODUCT CATEGORY:

Feed Additive

THERAPEUTIC GROUP:

Nutritional

PHARMACOTHERAPEUTIC GROUP:

Herbal extracts

COMPOSITION:

Garlic extract.....	120000 mg/L
Peppermint Extract.....	4000 mg/L
Lemon Peel Extract.....	100 mg/L
Cinnamon.....	40 mg/L
Ginger.....	40 mg/L

Withdrawal Times:

None

Storage:

Dark, dry and bellow 25 C. Do not freeze.

TARGET SPECIES:

Poultry, horse, cattle, camel, sheep, goat, pig

DIRECTIONS FOR USE:

ANIGARLIMINT SUPER oral solution is a feed additive, garlic and other herbal extracts in its content

- have an antioxidant effect and can be used for prevention of free radicals damaging the cell walls.
- Mint in its content deodorizes the smell of garlic while increasing appetite and facilitating digestion.
- can be used as an aromatic and appetizer in poultry.
- shows antimicrobial and anti-inflammatory effects,
- supports the immune system.
- supports the feed conversion-digestion rates, appetite, bodyweight, performance, accelerate growth
- can be used to eliminate the heat stress, prevent the loses of animals

DOSAGE:

Broiler and layers: 200-250 mL/ton drinking water Cattle, camels and horses: 20-30 mL/animal Calves, sheep, goats and pigs: 5-10 mL/animal

PRESENTATION: 1 L

HAYAT ROKHAN LTD

ANITONIC

Solution for Oral

PRODUCT CATEGORY:

Feed Additive

THERAPEUTIC GROUP:

Nutritional

PHARMACOTHERAPEUTIC GROUP:

Vitamin & Yeast

COMPOSITION:

Vitamin B1.....	250 mg/L
Vitamin B2.....	0.045 mg/L
Niacin.....	2500 mg/L
Inositol.....	2000 mg/L
Choline Chloride.....	100000 mg/L
L-carnitine.....	25000 mg/L
Betaine hydrochloride.....	10000 mg/L
Calcium-D-pantothenate.....	250 mg/L
Sorbitol.....	100000 mg/L
Active yeast extract (Saccharomyces cerevisiae).....	10.5x10 ⁷ CFU/L

Withdrawal Times:

None

Storage:

Dark, dry and bellow 25 C. Do not freeze.

TARGET SPECIES:

Poultry, pig

DIRECTIONS FOR USE:

ANITONIC is a feed additive combined of vitamins, pro vitamins and yeast. Strategic vitamin supplementation is aimed at protection of liver. It is used to support the immune function, animal growth, productivity and bioavailability of feed as well as body resistance against heavy stress conditions such as transplant, beak cut, vaccination and seasonal changes.

DOSAGE:

Poultry: 1-2 mL/4 L drinking water

Pigs: 1-2 mL/4 L drinking water

PRESENTATION: 1 L



ANIMIN

Solution for Oral

PRODUCT CATEGORY:

Feed Additive

THERAPEUTIC GROUP:

Nutritional

PHARMACOTHERAPEUTIC GROUP:

Mineral Combination

COMPOSITION:

Ferrous chloride (Fe).....	8000 mg/L
Manganous chloride (Mn).....	12500 mg/L
Zinc chloride (Zn).....	7000 mg/L
Cupric chloride (Cu).....	3500 mg/L
Calcium chloride (Ca).....	30000 mg/L
Phosphoric acid.....	30000 mg/L

Withdrawal Times:

None

Storage:

Dark, dry and bellow 25 C. Do not freeze.

TARGET SPECIES:

Poultry, ruminant

DIRECTIONS FOR USE:

ANIMIN oral solution is a nutritional feed additive which is advised

- in conditions of mineral (Fe, Mn, Cu, Zn and Ca) and phosphoric acid deficiencies
- to support the following cases: Perosis, cannibalism, cage fatigue, pica, stress, alopecia, dehydration, non-specific diarrhea, bone disorders, anemia, and metabolic defects

DOSAGE:

Poultry: 500 mL-1 L/1000 LI drinking water/day, for 3-5 days

Cattle, camels and horses: 20-30 mL/1 L drinking water/day, for 3-5 days

Calves, sheep and goats: 5 mL-10 mL/1 L drinking water/day, for 3-5 days

PRESENTATION: 1 L

HAYAT ROKHAN LTD

ANIMIN SUPER

Solution For Oral

PRODUCT CATEGORY:

Feed Additive

THERAPEUTIC GROUP:

Nutritional

PHARMACOTHERAPEUTIC GROUP:

Mineral Combination

COMPOSITION:

Ferrous chloride (Fe).....	1583 mg/L
Manganous chloride (Mn).....	582 mg/L
Zinc chloride (Zn).....	623 mg/L
Cupric chloride (Cu).....	238 mg/L
Cobalt chloride (Co).....	228 mg/L
Potassium chloride (K).....	2030 mg/L
Magnesium chloride (Mg).....	26940 mg/L
Sodium chloride (Na).....	4000 mg/L
Calcium chloride (Ca).....	30000 mg/L
Phosphoric acid.....	30000 mg/L

Withdrawal Times:

None

Storage:

Dark, dry and bellow 25 C. Do not freeze.

TARGET SPECIES:

Poultry, ruminant, horse

DIRECTIONS FOR USE:

ANIMIN SUPER oral solution is a nutritional feed additive which is advised

- in conditions of mineral (Fe, Mn, Co, Cu, Zn, K, Na, Mg and Ca) and phosphoric acid deficiencies
- to support the following cases: Perosis, cannibalism, cage fatigue, pica, stress, alopecia, dehydration, non-specific diarrhea, bone disorders, anemia, and metabolic defects

DOSAGE:

Poultry: 2.5-5 L/1000 L drinking water/day, for 3-5 days Cattle, camels and horses: 20-30 mL/1 L drinking water/day, for 3-5 days Calves, sheep and goats: 10-20 mL/1 L drinking water, for 3- 5 days

PRESENTATION: 1 L



ANIRES P

Solution For Oral

PRODUCT CATEGORY:

Feed Additive

THERAPEUTIC GROUP:

Nutritional

PHARMACOTHERAPEUTIC GROUP:

Essential Oils (EOs)

COMPOSITION:

Eucalyptus oil.....	35000 mg/L
Mint oil.....	14000 mg/L
Thyme oil.....	7000 mg/L
Menthol.....	21000 mg/L

Withdrawal Times:

None

Storage:

Dark, dry and bellow 25 C. Do not freeze.

TARGET SPECIES:

Poultry, ruminant, horse, pig

DIRECTIONS FOR USE:

ANIRES P is a herbal feed additive product which protects the upper airways and have an expectorant effect on the mucus membranes of nose, sinuses and trachea. It is advised to use during periods of disadvantageous environmental conditions (high temperature, high pollination) or post-vaccination to alleviate the undesirable effects.

DOSAGE:

Poultry: 250-300 mL/1000 L drinking water

Pigs, calves: 2,5-3 mL/10 L drinking water

Horses: 20-25 mL/animal

PRESENTATION: 1 L

HAYAT ROKHAN LTD

ANIVIT AD3ECK

Solution For Oral

PRODUCT CATEGORY:

Feed Additive

THERAPEUTIC GROUP:

Nutritional

PHARMACOTHERAPEUTIC GROUP:

Vitamin

COMPOSITION:

Vitamin A.....10.000.000 IU/L
Vitamin D3.....1.000.000 IU/L
Vitamin E.....8500 mg/L
Vitamin C.....8000 mg/L
Vitamin K.....5000 mg/L

Withdrawal Times:

None

Storage:

Dark, dry and bellow 25 C. Do not freeze.

TARGET SPECIES:

Poultry, ruminant, horse

DIRECTIONS FOR USE:

ANIVIT AD3ECK With its content of vitamin A, vitamin D3, vitamin E, vitamin C and vitamin K, it is used in case of vitamin deficiencies. It is also used as a supportive for

- bone and skeletal system disorders
- stress conditions (due to transport, beak cutting, vaccination, temperature, feed and climate changes)
- immune system with its antioxidant properties
- body growth, reproductive performance and feed conversion rate (FCR)
- blood clotting

DOSAGE:

Poultry:

Broiler: 500 mL- 1L/1000 L drinking water

Layers and breeders: 1 -1,5 L/1000 L drinking water

Calves and foals: 5 mL/animal, through mixing with drinking water

Cattle, camels and horses: 10 mL/animal, through mixing with drinking water

PRESENTATION: 1 L



ANIVIT AMINO

Solution For Oral

PRODUCT CATEGORY:

Feed Additive

THERAPEUTIC GROUP:

Nutritional

PHARMACOTHERAPEUTIC GROUP:

Vitamin & Amino acid

COMPOSITION:

Vitamin A.....	10.000.000 IU/L	L-Lysine.....	3000 mg/L
Vitamin D3.....	1.000.000 IU/L	L-Arginine.....	600 mg/L
Vitamin E.....	10000 mg/L	L-Leucine.....	600 mg/L
Vitamin C.....	25000 mg/L	L-Valine.....	600 mg/L
Vitamin B1.....	2500 mg/L	L- Tyrosine.....	200 mg/L
Vitamin B2.....	1250 mg/L	L-Isoleucine.....	200 mg/L
Vitamin B6.....	500 mg/L	L-Proline.....	800 mg/L
Vitamin B12.....	5 mg/L	L-Threonine.....	500 mg/L
Vitamin K3.....	500 mg/L	L-Tryptophan.....	20 mg/L
D-Biotin.....	15 mg/L	L-Glutamic acid.....	4.000 mg/L
Calcium D-Pantothenate.....	2500 mg/L		
Choline chloride.....	70000 mg/L		
DL-Methionine.....	550 mg/L		

Withdrawal Times:

None

Storage:

Dark, dry and bellow 25 C. Do not freeze.

TARGET SPECIES:

Poultry, ruminant, horse, pet, pig

DIRECTIONS FOR USE:

ANIVIT AMINO Solution for oral recommended to be used as a supportive supplement in cases of dehydration, electrolyte, vitamin imbalance, hyperproteinemia which can occur in diarrhea, shock and exhaustion. Recommended to be used as a preventative supplement in cases of breeding losses, nutritional deficiencies cases, fatty liver syndrome and nerve disorders. Can also be used during period of convalescence.

DOSAGE:

Poultry: 250-500 mL/ton drinking water, for 3-5 days
Horse: 20-30 mL/day Calves: 10-20 mL/day
Pigs: 5-10 mL Lambs: 2-4 mL/day Cats and dogs: 1-2 mL/day

PRESENTATION: 1 L

ANIVIT CARLYMES

Solution For Oral

PRODUCT CATEGORY:

Feed Additive

THERAPEUTIC GROUP:

Nutritional

PHARMACOTHERAPEUTIC GROUP:

Amino acid

COMPOSITION:

L-Carnitine.....	50000 mg/L
L-Lysine.....	1000 mg/L
L-Methionine.....	10000 mg/L
VitaminC.....	50000 mg/L
Sorbitol.....	250000 mg/L

Withdrawal Times:

None

Storage:

Dark, dry and bellow 25 C. Do not freeze.

TARGET SPECIES:

Poultry, ruminant

DIRECTIONS FOR USE:

ANIVIT CARLYMES is used

- to increase efficiency, to improve yield (Carcass quality etc.), to increase the feed conversion rate and to support the immune mechanism,
- in nutritional deficiencies seen during bacterial and viral diseases,
- during convalescence period of diseases,
- to increase specific antibody production,
- in stress conditions,
- in fatty liver and kidney syndrome and fatty liverhemorrhagic syndrome,
- as a supporting agent in cases of ammonia detoxification

DOSAGE:

Poultry: 1-2 L/ 1000 L drinking water/day

Calves and lambs: 10-15 mL/animal

Pigs: 5-10 mL/animal

PRESENTATION: 1 L